



*Caroline Gerhart,
Registered Dental Assistant*



MARK F. HARDISON, D.D.S. & JEFF WARDEN, D.D.S., M.D.



Our goal is to provide you with a safe, pleasant and effective anesthetic. In order to do this it is imperative that we have your full cooperation. Please feel free to call us at **615-893-7736** (Murfreesboro office) or **615-462-7987** (Smyrna office) concerning questions regarding your surgery or anesthetic.



MIDDLE TENNESSEE ORAL & IMPLANT SURGERY

Compassion. Availability. Excellence.

IMPORTANT INSTRUCTIONS

Regarding Your Anesthesia



MIDDLE TENNESSEE ORAL & IMPLANT SURGERY

1725 MEDICAL CENTER PKWY., STE. 100
MURFREESBORO, TENNESSEE 37129
P: 615-893-7736 | F: 615-898-1771

515 STONECREST PKWY., STE. 240
SMYRNA, TENNESSEE 37167
P: 615-462-7987 | F: 615-625-3405

WWW.MTOIS.COM



*Shelby Daniel,
Financial Coordinator*



You will always be given local anesthesia for your surgery, but you may choose any of those listed below as a supplement. Each choice requires different preparation on your part, and for your safety it is important that you read and follow the instructions carefully. If you are unclear about anything, please ask your doctor.

For all surgeries, please wear comfortable, loose-fitting clothing. Tops/shirts should have sleeves that are easily drawn up above the elbow. Females should remove nail polish before surgery, and apply as little makeup as possible.

LOCAL ANESTHESIA

Will produce a numb feeling in the area being operated on and a feeling of pressure during surgery. You will be awake and recall the surgery, but there should be no significant discomfort.

1. Have a light meal a few hours prior to surgery.
2. For more extensive procedures you may wish to have someone drive you home.
3. Plan to rest for a few hours after surgery.

NITROUS OXIDE

is also known as “laughing gas.” You will be relaxed and somewhat less aware of your surroundings, but will recall most of the surgical event. Nitrous oxide is generally used in conjunction with local anesthesia, but may also be used to supplement the anesthetic choices below.

1. You may have a light meal four (4) hours prior to surgery.
2. It is best to have someone drive you home.
3. Plan to rest for the remainder of the day.

INTRAVENOUS ANESTHESIA

Both of the anesthetics below include local anesthesia (although general anesthesia does not require its use.)

INTRAVENOUS SEDATION:

Medications are given through a vein in your arm or hand, which will cause total relaxation and, although you will not actually be unconscious, there will be very little recall (if any) of the events surrounding surgery.

IMPORTANT INSTRUCTIONS REGARDING YOUR ANESTHESIA

1. Do not eat or drink anything (including water) for six (6) hours prior to surgery. However, it is important that you take any regular medications (high blood pressure, antibiotics, etc.) or any pre-medication prescription that we have provided, using only a small sip of water.
2. For morning appointments, skip breakfast.
3. For afternoon appointments, eat a light breakfast seven (7) hours before your appointment and skip lunch.
4. Take any regular medications with only enough water to get the pill down.
5. You **MUST** have someone drive you home and be onsite for your entire office procedure.
6. Plan to rest for the remainder of the day. Do not operate power tools, machinery, etc., for 24 hours after surgery.

GENERAL ANESTHESIA:

Medications are given through a vein which will result in total loss of consciousness, complete lack of recall of the event and usually a longer recovery time. General anesthesia has an excellent safety record as an office procedure, but may, if desired, be provided in a hospital setting. (Your health insurance may not cover you unless there is a bona fide medical reason for hospitalization.)

The same instructions offered above for intravenous sedation apply for general anesthesia.

